


PHARMACY POLICY – 5.01.520

Antidepressants: Pharmacy Medical Necessity Criteria for Brands

Effective Date:	Jan. 1, 2025	RELATED MEDICAL POLICIES:
Last Revised:	Dec. 10, 2024	5.01.521 Pharmacologic Treatment of Neuropathy, Fibromyalgia, and Seizure Disorders
Replaces:	N/A	5.01.605 Medical Necessity Criteria for Pharmacy Edits

Select a hyperlink below to be directed to that section.

[POLICY CRITERIA](#) | [CODING](#) | [RELATED INFORMATION](#)
[EVIDENCE REVIEW](#) | [REFERENCES](#) | [HISTORY](#)

 Clicking this icon returns you to the hyperlinks menu above.

Introduction

Drugs are grouped into classes. Drugs can be in the same class because they work in the same way, have a similar chemical structure, or are used for the same purpose. There are a number of drug classes for antidepressant drugs. These include monoamine oxidase inhibitors (MAOIs), tricyclic antidepressants (TCAs), selective serotonin reuptake inhibitors (SSRIs), and serotonin-norepinephrine reuptake inhibitors (SNRIs). MAOIs and TCAs were the first drugs developed in the 1950s and are considered first-generation antidepressants. Because antidepressants like SSRIs and SNRIs were developed later they are considered second-generation antidepressants. Like most drugs, second-generation antidepressants come in both brand and generic form. The active ingredients in generic drugs are chemically identical to the active ingredients in brand name drugs. Because generic drugs have the same active ingredients as brand name drugs, they are usually tried first. This policy describes when brand-name second-generation antidepressants may be considered medically necessary.

Note: The Introduction section is for your general knowledge and is not to be taken as policy coverage criteria. The rest of the policy uses specific words and concepts familiar to medical professionals. It is intended for providers. A provider can be a person, such as a doctor, nurse, psychologist, or dentist. A provider also can be a place where medical care is given, like a hospital, clinic, or lab. This policy informs them about when a service may be covered.

Policy Coverage Criteria

Condition & Drug	Medical Necessity
<p>Anxiety</p> <ul style="list-style-type: none"> • Aplenzin ER (bupropion hydrobromide ER) • Bupropion XL 450mg tablet • Celexa (citalopram) • Citalopram capsule • Cymbalta (duloxetine) • Desvenlafaxine ER • Drizalma Sprinkle (duloxetine DR) • Effexor XR (venlafaxine ER) • Fetzima (levomilnacipran) • Forfivo XL (bupropion ER) • Lexapro (escitalopram) • Paxil (paroxetine) • Paxil CR (paroxetine ER) • Pristiq (desvenlafaxine) • Prozac (fluoxetine) • Remeron (mirtazapine) • Sertraline capsule • Symbyax (olanzapine and fluoxetine) • Trintellix (vortioxetine) • Venlafaxine Besylate ER • Viibryd (vilazodone) • Wellbutrin SR (bupropion SR) • Wellbutrin XL (bupropion ER) • Zoloft (sertraline) 	<p>Brand selective serotonin reuptake inhibitor (SSRI), serotonin/norepinephrine reuptake inhibitor (SNRI), and any second-generation antidepressant, when used to treat anxiety, may be considered medically necessary when:</p> <ul style="list-style-type: none"> • The individual has had a trial and failure of two generic SSRIs (e.g., citalopram, escitalopram, fluoxetine, paroxetine, sertraline) <p>OR</p> <ul style="list-style-type: none"> • Has had a trial and failure of two generic SNRIs (e.g., desvenlafaxine, duloxetine, venlafaxine) <p>OR</p> <ul style="list-style-type: none"> • Has had a trial and failure of one generic SSRI and one generic SNRI <p>Note: For diagnosis of diabetic peripheral neuropathy, chronic musculoskeletal pain or fibromyalgia, see Related Policies.</p>
<p>Depression</p>	<p>Brand selective serotonin reuptake inhibitor (SSRI), serotonin/norepinephrine reuptake inhibitor (SNRI), and any</p>



Condition & Drug	Medical Necessity
<ul style="list-style-type: none"> • Aplenzin ER (bupropion hydrobromide ER) • Auvelity (dextromethorphan and bupropion) • Bupropion XL 450mg tablet • Celexa (citalopram) • Citalopram capsule • Cymbalta (duloxetine) • Desvenlafaxine ER • Drizalma Sprinkle (duloxetine DR) • Effexor XR (venlafaxine ER) • Exxua (gepirone) • Fetzima (levomilnacipran) • Forfivo XL (bupropion ER) • Lexapro (escitalopram) • Paxil (paroxetine) • Paxil CR (paroxetine ER) • Pristiq (desvenlafaxine) • Prozac (fluoxetine) • Remeron (mirtazapine) • Sertraline capsule • Symbyax (olanzapine and fluoxetine) • Trintellix (vortioxetine) • Venlafaxine Besylate ER • Viibryd (vilazodone) • Wellbutrin SR (bupropion SR) • Wellbutrin XL (bupropion ER) • Zoloft (sertraline) 	<p>second-generation antidepressant, when used to treat depression, may be considered medically necessary when:</p> <ul style="list-style-type: none"> • The individual has had a trial and failure of two generic second-generation antidepressants (e.g., fluoxetine, sertraline, venlafaxine, bupropion, mirtazapine) <p>Auvelity (dextromethorphan and bupropion) when used to treat depression, may be considered medically necessary when:</p> <ul style="list-style-type: none"> • The individual has had a trial and failure of two generic second-generation antidepressants (e.g., fluoxetine, sertraline, venlafaxine, bupropion, mirtazapine) <p>Exxua (gepirone) when used to treat depression, may be considered medically necessary when:</p> <ul style="list-style-type: none"> • The individual has had a trial and failure of two generic second-generation antidepressants (e.g., fluoxetine, sertraline, venlafaxine, bupropion, mirtazapine) <p>Note: For diagnosis of diabetic peripheral neuropathy, chronic musculoskeletal pain or fibromyalgia, see Related Policies.</p>

Drug	Not Medically Necessary
As listed	All other uses of the medications listed in this policy are considered not medically necessary.



Drug	Investigational
As listed	The medications listed in this policy are subject to the product's U.S. Food and Drug Administration (FDA) dosage and administration prescribing information.

Length of Approval	
Approval	Criteria
Initial authorization	The medications listed in this policy may be approved up to 3 years.
Re-authorization criteria	Future re-authorization of the medications listed in this policy may be approved up to 3 years as long as the medical necessity criteria are met and chart notes demonstrate that the individual continues to show a positive clinical response to therapy.

Documentation Requirements
<p>The individual's medical records submitted for review for all conditions should document that medical necessity criteria are met. The record should include the following:</p> <ul style="list-style-type: none"> Office visit notes that contain the diagnosis, relevant history, physical evaluation and medication history

Coding

N/A

Related Information



Benefit Application

This policy applies to all pharmacy benefit contracts that include Pharmacy Prior Authorization Edits.

This policy is managed through the Pharmacy benefit.

Evidence Review

Description

Pathophysiology of Depression

While the pathology of depression is far from completely understood, it is apparent that both heredity and environmental factors play a part. Genetic microarray techniques are being used to identify candidate genes, with the hope of developing a pharmacogenomic approach to predicting which drugs will be most effective in each individual. However, this knowledge is still in its infancy, and its practical application will be some time in the future. For now, practitioners must continue using empiric approaches to treatment, both pharmacologic and otherwise.

Disease Burden

More than 18 million Americans suffer from depression. Over 15% will experience at least one major depressive episode during their lifetimes. Exact prevalence rates are difficult to determine because of the extent to which depression goes unreported. It occurs twice as frequently in women as in men. A meta-analysis showed that depressed persons have a 1.5-2 fold increased risk of mortality.

In 2000, the economic burden of depression in the US was estimated to be \$83.1 billion. Indirect costs include related mortality and morbidity, as well as significant amounts of absenteeism and presenteeism that can impact workplace productivity. Quality of life of individuals and those around them also suffer.



Pharmacotherapy

An overview of the various treatment modalities used in depression was provided in Lancet by Ebmeier et al. Treatments include a variety of cognitive behavioral approaches, psychotherapy, treatment with antidepressant medications and in severely resistant cases, electroconvulsive therapy. Of the nonpharmacologic treatment modalities, cognitive behavioral approaches have the best supporting evidence. Drugs used to treat depression include selective serotonin reuptake inhibitors (SSRIs), tricyclic antidepressants (TCAs), monoamine oxidase inhibitors (MAOI) and several agents with combinations of serotonergic, noradrenergic and dopaminergic activity.

This policy applies to the following medications:

- Single Source Brand (SSB) selective serotonin reuptake inhibitor (SSRI), serotonin/norepinephrine reuptake inhibitor (SNRI) and any other SSB second-generation antidepressants (antidepressants other than tricyclic and MAOI agents).

Summary of Evidence

Current evidence from head-to-head comparative trials of second-generation antidepressants, systematic reviews, and meta-analyses indicates these agents are of comparable efficacy and effectiveness as measured by HAM-D or MADRS response (>50% improvement), though one study reported a statistically superior but modest improvement in MADRS score with escitalopram compared to citalopram. Overall, the data support for a meaningful difference is not compelling.

Evidence from four small comparative effectiveness trials failed to provide compelling evidence of the superiority of escitalopram. In the one study that showed a statistically significant difference in the primary endpoint (change from baseline in MADRS score), the effect size was modest, and p value was barely significant.

Two longer-term (≥6 month) efficacy and safety studies of duloxetine (Cymbalta) duloxetine have been published, and the manufacturer supplied data on two unpublished trials. Although not compelling, evidence now supports the relative safety of longer-term (6 month to 1 year) use of duloxetine for the treatment of major depressive disorder. Other evidence also shows comparable efficacy with venlafaxine and comparable onset of effect with escitalopram.

A meta-analysis of 117 small randomized controlled trials including 25,928 individuals compared 12 “new generation” antidepressants for efficacy and tolerability. The authors concluded that



escitalopram and sertraline offered the best combination of efficacy and individual acceptability. Of the two, they felt that sertraline might be the best choice when starting treatment, because it has the best balance between efficacy, safety and cost. Although well-designed, this study is limited by the size and heterogeneity of the individual trials that were included. In particular, the evidence supporting superiority of escitalopram over citalopram is based on 5 small trials, all of which were manufacturer-sponsored. There is no clinical reason to expect a meaningful difference between these two agents, assuming that the doses of the S-isomer are equal.

A similar meta-analysis of 203 studies yielded no substantial differences among agents. The authors concluded that the body of existing evidence does not favor selection of one particular antidepressant over the others based on efficacy or effectiveness.

US guidelines for the treatment of adults with depression from the American Psychiatric Association recommend use of antidepressants as preferred initial treatment or as a part of a preferred initial treatment regimen for most individuals with any level of severity of MDD. Initial choice of medication should consider anticipated side effects, safety or tolerability of side effects for individual individuals, individual preference, quantity and quality of clinical trial data, and cost. Based on these factors, the APA indicates SSRIs, desipramine, nortriptyline, bupropion, venlafaxine, and mirtazapine are likely to be effective for most individuals.

2009 Update

A literature search was performed for March 2009 through December 2009. No published randomized studies were found that would change the policy statements.

2011 Update

Updated to incorporate reference to newly US Food and Drug Administration (FDA-approved indication for Cymbalta in chronic musculoskeletal pain). No other significant updates to the literature were found.

2012 Update

Updated to allow for recent availability of generic escitalopram; thus, trial of generic citalopram is no longer a specific requirement for access to Lexapro. No other significant updates to the literature were found.



2014 Update

Updated to include newly available generic duloxetine and Khedezla, a new venlafaxine extended release product. No other significant changes to the literature were found at this time.

2015 Update

Expanded the indication of Major Depressive Disorder to Depressive Disorders; Expanded the indication of Generalized Anxiety Disorder to Anxiety Disorders. Updated these new expanded indications with separate criteria: For individuals with Depressive Disorders, trial and failure of at least two generically available second generation antidepressants; or individuals with Anxiety Disorders, trial and failure of at least two generically available SSRIs, or at least one generically available SSRI and one generically available SNRI.

2016 Update

A literature search was performed for April 1, 2015 through December 6, 2016. No published randomized studies were found that would change existing policy statements.

2017 Update

A literature search was performed for July 1, 2016 through November 2, 2017. No published studies were found that would change existing policy statements.

2019 Update

A literature search was performed for December 1, 2018 through November 30, 2019. No published studies were found that would change existing policy statements.



2020 Update

No changes to policy statement. Combined depression and anxiety criteria into one table. Added a table documenting that all other uses of brand SSRI, SNRI, and any second- generation antidepressant for this policy are considered not medically necessary. Added a Length of Approval and Documentation Requirements table to policy.

2021 Update

A literature search was performed for July 1, 2020 through June 30, 2021. No published studies were found that would change existing policy statements.

2022 Update

Reviewed UpToDate on “Unipolar major depression in adults: Choosing initial treatment”. Evidence presented continues to support initial treatment with an SSRI or SNRI rather than other antidepressants. No new information was identified that would require changes to this policy.

2023 Update

No new information was identified that would require changes to this policy.

2024 Update

Added Exxua (gepirone) coverage criteria for the treatment of depression.

References

1. Rush, AJ, Pincus HA, First MB et al. Handbook of Psychiatric Measures, First Edition, pp100-102. 2000; Washington, DC, American Psychiatric Association Press. Rush AJ, op. cit., pp526-540.
2. Gartlehner G, Hansen RA, Kahwati L. Drug Class Review on Second Generation Antidepressants. 2011. Available at <http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0009808/> Accessed February 23, 2024.



3. Lepola UM, Loft H, Reines EH. Escitalopram (10-20 mg/day) is effective and well tolerated in a placebo-controlled study in depression in primary care. *Int Clin Psychopharmacol.* 2003;18(4):211-217.
4. Burke WJ, Gergel I, Bose A. Fixed-dose trial of the single isomer SSRI escitalopram in depressed outpatients. *J Clin Psychiatry.* 2002;63(4):331-336.
5. Colonna L, Andersen HF, Reines EH. A randomized, double-blind, 24-week study of escitalopram (10 mg/day) versus citalopram (20 mg/day) in primary care patients with major depressive disorder. *Curr Med Res Opin.* 2005;21(10):1659-1668.
6. Moore N, Verdoux H, Fantino B. Prospective, multicentre, randomized, double-blind study of the efficacy of escitalopram versus citalopram in outpatient treatment of major depressive disorder. *Int Clin Psychopharmacol.* 2005;20(3):131-137.
7. Sechter D, Troy S, Paternetti S, Boyer P. A double-blind comparison of sertraline and fluoxetine in the treatment of major depressive episode in outpatients. *Eur Psychiatry.* 1999;14(1):41-48.
8. Fava M, Hoog SL, Judge RA, Kopp JB, Nilsson ME, Gonzales JS. Acute efficacy of fluoxetine versus sertraline and paroxetine in major depressive disorder including effects of baseline insomnia. *J Clin Psychopharmacol.* 2002;22(2):137-147.
9. Bennie EH, Mullin JM, Martindale JJ. A double-blind multicenter trial comparing sertraline and fluoxetine in outpatients with major depression. *J Clin Psychiatry.* 1995;56(6):229-237.
10. Thompson C. Management of depression in real-life settings: knowledge gained from large scale clinical trials. *Int Clin Psychopharmacol.* 1994;9 Suppl 3:21-25.
11. Newhouse PA, Krishnan KR, Doraiswamy PM, Richter EM, Batzar ED, Clary CM. A double-blind comparison of sertraline and fluoxetine in depressed elderly outpatients. *J Clin Psychiatry.* 2000;61(8):559-568.
12. Chouinard G, Saxena B, Belanger MC, Ravindran A, Bakish D, Beauclair L et al. A Canadian multicenter, double-blind study of paroxetine and fluoxetine in major depressive disorder. *J Affect Disord.* 1999;54(1-2):39-48.
13. De Wilde J, Spiers R, Mertens C, Bartholome F, Schotte G, Leyman S. A double-blind, comparative, multicentre study comparing paroxetine with fluoxetine in depressed patients. *Acta Psychiatr Scand.* 1993;87(2):141-145.
14. Schone W, Ludwig M. A double-blind study of paroxetine compared with fluoxetine in geriatric patients with major depression. *J Clin Psychopharmacol.* 1993;13(6 Suppl 2):34S-39S.
15. Fava M, Amsterdam JD, Deltito JA, Salzman C, Schwaller M, Dunner DL. A double-blind study of paroxetine, fluoxetine, and placebo in outpatients with major depression. *Ann Clin Psychiatry.* 1998;10(4):145-150.
16. Fava M, Rosenbaum JF, Hoog SL, Tepner RG, Kopp JB, Nilsson ME. Fluoxetine versus sertraline and paroxetine in major depression: tolerability and efficacy in anxious depression. *J Affect Disord.* 2000;59(2):119-126.
17. Cassano GB, Puca F, Scapicchio PL, Trabucchi M. Paroxetine and fluoxetine effects on mood and cognitive functions in depressed nondemented elderly patients. *J Clin Psychiatry.* 2002;63(5):396-402.
18. Gagiano CA. A double blind comparison of paroxetine and fluoxetine in patients with major depression. *B J Clin Res.* 1993;4:145-152.
19. Costa e Silva J. Randomized, double-blind comparison of venlafaxine and fluoxetine in outpatients with major depression. *J Clin Psychiatry.* 1998;59(7):352-357.
20. De Nayer A, Geerts S, Ruelens L, Schittecatte M, De Bleeker E, Van Eeckhoutte I et al. Venlafaxine compared with fluoxetine in outpatients with depression and concomitant anxiety. *Int J Neuropsychopharmacol.* 2002;5(2):115-120.
21. Rudolph RL, Feiger AD. A double-blind, randomized, placebo-controlled trial of once-daily venlafaxine extended release (XR) and fluoxetine for the treatment of depression. *J Affect Disord.* 1999;56(2-3):171-181.
22. Silverstone PH, Ravindran A. Once-daily venlafaxine extended release (XR) compared with fluoxetine in outpatients with depression and anxiety. Venlafaxine XR 360 Study Group. *J Clin Psychiatry.* 1999;60(1):22-28.
23. Alves C, Cachola I, Brandao J. Efficacy and tolerability of venlafaxine and fluoxetine in outpatients with major depression. *Primary Care Psychiatry.* 1999;5(2):57-63.



24. Dierick M, Ravizza L, Realini R, Martin A. A double-blind comparison of venlafaxine and fluoxetine for treatment of major depression in outpatients. *Prog Neuropsychopharmacol Biol Psychiatry*. 1996;20(1):57-71.
25. Tylee A, Beaumont G, Bowden MW, Reynolds A. A double-blind, randomized, 12-week comparison study of the safety and efficacy of venlafaxine and fluoxetine in moderate to severe depression in general practice. *Primary Care Psychiatry*. 1997;3:51-58.
26. Montgomery SA. Comparative efficacy and tolerability of escitalopram oxalate versus venlafaxine XR. Data on file, Forest Labs; 2004.
27. Bielski RJ, Ventura D, Chang CC. A double-blind comparison of escitalopram and venlafaxine extended release in the treatment of major depressive disorder. *J Clin Psychiatry*. 2004;65(9):1190-1196.
28. Schatzberg AF, Kremer C, Rodrigues HE, Murphy GMJ. Double-blind, randomized comparison of mirtazapine and paroxetine in elderly depressed patients. *Am J Geriatr Psychiatry*. 2002;10(5):541-550.
29. Benkert O, Szegedi A, Kohnen R. Mirtazapine compared with paroxetine in major depression. *J Clin Psychiatry*. 2000;61(9):656-663.
30. Behnke K, Sogaard J, Martin S, Bauml J, Ravindran AV, Agren H et al. Mirtazapine orally disintegrating tablet versus sertraline: a prospective onset of action study. *J Clin Psychopharmacol*. 2003;23(4):358-364.
31. Hong CJ, Hu WH, Chen CC, Hsiao CC, Tsai SJ, Ruwe FJ. A double-blind, randomized, group-comparative study of the tolerability and efficacy of 6 weeks' treatment with mirtazapine or fluoxetine in depressed Chinese patients. *J Clin Psychiatry*. 2003;64(8):921-926.
32. Feighner JP, Gardner EA, Johnston JA, Batey SR, Khayrallah MA, Ascher JA et al. Double-blind comparison of bupropion and fluoxetine in depressed outpatients. *J Clin Psychiatry*. 1991;52(8):329-335.
33. Coleman CC, King BR, Bolden-Watson C, Book MJ, Segraves RT, Richard N et al. A placebo-controlled comparison of the effects on sexual functioning of bupropion sustained release and fluoxetine. *Clin Ther*. 2001;23(7):1040-1058.
34. Weihs KL, Settle ECJ, Batey SR, Houser TL, Donahue RM, Ascher JA. Bupropion sustained release versus paroxetine for the treatment of depression in the elderly. *J Clin Psychiatry*. 2000;61(3):196-202.
35. Kavoussi RJ, Segraves RT, Hughes AR, Ascher JA, Johnston JA. Double-blind comparison of bupropion sustained release and sertraline in depressed outpatients. *J Clin Psychiatry*. 1997;58(12):532-537.
36. Croft H, Settle EJ, Houser T, Batey SR, Donahue RM, Ascher JA. A placebo-controlled comparison of the antidepressant efficacy and effects on sexual functioning of sustained release bupropion and sertraline. *Clin Ther*. 1999;21(4):643-658.
37. Coleman CC, Cunningham LA, Foster VJ, Batey SR, Donahue RM, Houser TL et al. Sexual dysfunction associated with the treatment of depression: a placebo-controlled comparison of bupropion sustained release and sertraline treatment. *Ann Clin Psychiatry*. 1999;11(4):205-215.
38. Nieuwstraten CE, Dolovich LR. Bupropion versus selective serotonin-reuptake inhibitors for treatment of depression. *Ann Pharmacother*. 2001;35(12):1608-1613.
39. Whittington CJ, Kendall T, Fonagy P, Cottrell D, Cotgrove A, Boddington E. Selective serotonin reuptake inhibitors in childhood depression: systematic review of published versus unpublished data. *Lancet*. 2004;363(9418):1341-1345.
40. Ekselius L, von Knorring L, Eberhard G. A double-blind multicenter trial comparing sertraline and citalopram in patients with major depression treated in general practice. *Int Clin Psychopharmacol*. 1997;12(6):323-331.
41. Sechter D, Troy S, Paternetti S, Boyer P. A double-blind comparison of sertraline and fluoxetine in the treatment of major depressive episode in outpatients. *Eur Psychiatry*. 1999;14(1):41-48.
42. Kroenke K, West SL, Swindle R, Gilson A, Eckert GJ, Dolor R et al. Similar effectiveness of paroxetine, fluoxetine, and sertraline in primary care: a randomized trial. *JAMA*. 2001;286(23):2947-2955.
43. Raskin J, Goldstein DJ, Mallinckrodt CH, Ferguson MB. Duloxetine in the long-term treatment of major depressive disorder. *J Clin Psychiatry*. 2003;64(10):1237-44.



44. Perahia, DG, Wang F, Mallinckrodt CH et al. Duloxetine in the treatment of major depressive disorder: a placebo- and paroxetine-controlled trial. *Eur Psychiatry*. May 10, 2006;[Epub ahead of print].
45. Goldstein DJ, Lu Y, Detke MJ et al. Duloxetine in the treatment of depression : a double-blind placebo-controlled comparison with paroxetine. *J Clin Psychopharmacol*. 2004;24(4):389-399.
46. Mackay FJ, Dunn NR, Wilton LV, Pearce GL, Freemantle SN, Mann RD. A comparison of fluvoxamine, fluoxetine, sertraline and paroxetine examined by observational cohort studies. *Pharmacoepid Drug Safety*. 1997;6:235-246.
47. Segraves RT, Kavoussi R, Hughes AR, Batey SR, Johnston JA, Donahue R et al. Evaluation of sexual functioning in depressed outpatients: a double-blind comparison of sustained release bupropion and sertraline treatment. *J Clin Psychopharmacol*. 2000;20(2):122-128.
48. Clayton AH, Pradko JF, Croft HA, Montano CB, Leadbetter RA, Bolden-Watson C et al. Prevalence of sexual dysfunction among newer antidepressants. *J Clin Psychiatry*. 2002;63(4):357-366.
49. Aberg-Wistedt A, Agren H, Ekselius L, Bengtsson F, Akerblad AC. Sertraline versus paroxetine in major depression: clinical outcome after six months of continuous therapy. *J Clin Psychopharmacol*. 2000;20(6):645-652.
50. Nemeroff CB, Ninan PT, Ballenger J, Lydiard RB, Feighner J, Patterson WM et al. Double blind multicenter comparison of fluvoxamine versus sertraline in the treatment of depressed outpatients. 1995;3:163-169.
51. Feiger A, Kiev A, Shrivastava RK, Wisselink PG, Wilcox CX. Nefazodone versus sertraline in outpatients with major depression: focus on efficacy, tolerability, and effects on sexual function and satisfaction. *J Clin Psychiatry*. 1996;57(Suppl 2):53-62.
52. Fava M, Judge R, Hoog SL, Nilsson ME, Koke SC. Fluoxetine versus sertraline and paroxetine in major depressive disorder: changes in weight with long-term treatment. *J Clin Psychiatry*. 2000;61(11):863-867.
53. Michelson D, Amsterdam JD, Quitkin FM, Reimherr FW, Rosenbaum JF, Zajecka J et al. Changes in weight during a 1-year trial of fluoxetine. *Am J Psychiatry*. 1999;156(8):1170-1176.
54. Maina G, Albert U, Salvi V, Bogetto F. Weight gain during long-term treatment of obsessive-compulsive disorder: a prospective comparison between serotonin reuptake inhibitors. *J Clin Psychiatry*. 2004;65(10):1365-1371.
55. Croft H, Houser TL, Jamerson BD, Leadbetter R, Bolden-Watson C, Donahue R et al. Effect on body weight of bupropion sustained-release in patients with major depression treated for 52 weeks. *Clin Ther*. 2002;24(4):662-672.
56. Thase ME. Effects of venlafaxine on blood pressure: a meta-analysis of original data from 3744 depressed patients. *J Clin Psychiatry*. 1998;59(10):502-508.
57. Thase ME, Tran PV, Wiltse C, Pangallo BA, Mallinckrodt C, Detke MJ. Cardiovascular profile of duloxetine, a dual reuptake inhibitor of serotonin and norepinephrine. *J Clin Psychopharmacol*. 2005;25(2):132-140.
58. Buckley NA, McManus PR. Fatal toxicity of serotonergic and other antidepressant drugs: analysis of United Kingdom mortality data. *BMJ*. 2002;325(7376):1332-1333.
59. Liu BA, Mittmann N, Knowles SR, Shear NH. Hyponatremia and the syndrome of inappropriate secretion of antidiuretic hormone associated with the use of selective serotonin reuptake inhibitors: a review of spontaneous reports. *CMAJ*. 1996;155(5):519-527.
60. Stewart DE. Hepatic adverse reactions associated with nefazodone. *Can J Psychiatry*. 2002;47(4):375-377.
61. Fochtmann LJ, Gelenberg AJ. Guideline watch (2005): practice guideline for the treatment of patients with major depressive disorder, 2nd edition. Available at: https://psychiatryonline.org/pb/assets/raw/sitewide/practice_guidelines/guidelines/bipolar-watch.pdf Accessed February 23, 2024.
62. American Psychiatric Association. Practice guideline for the treatment of patients with major depressive disorder (revision). *Am J Psychiatry*. 2000;157:1-45.
63. American Psychiatric Association. Practice Guideline for the Treatment of Patients with Major Depressive Disorder. <http://focus.psychiatryonline.org/doi/abs/10.1176/foc.3.1.34?journalCode=foc> Accessed February 23, 2024.



64. National Institute for Clinical Excellence (NICE). Depression in adults: The treatment and management of depression in adults. Last updated June 2022. Available at: Overview | Depression in adults: treatment and management | Guidance | NICE Accessed February 23, 2024.
65. Barrett B, Byford S, Knapp M. Evidence of cost-effective treatments for depression: a systematic review. *J Affect Disord.* 2005;84(1):1-13.
66. Dunn JD, Cannon E, Mitchell MP, Curtiss FR. Utilization and drug cost outcomes of a step-therapy edit for generic antidepressants in an HMO in an integrated health system. *J Manag Care Pharm.* 2006; 12(4):294-302.
67. Kessler RC, Berglund P, Demler O et al. The epidemiology of major depressive disorder: results from the National Comorbidity Survey Replication (NCS-R). *National Comorbidity Survey Replication. JAMA.* 2003;289:3095-3105.
68. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, 4th Edition.* Washington, DC: American Psychiatric Association, 1994.
69. Cuijpers P, Smit F. Excess mortality in depression: a metaanalysis of community studies. *J Affect Disord* 2002; 72: 227–36, cited in Ebmeier KP, Donaghey C and Steele JD. Recent developments and current controversies in depression. *Lancet* 2006;367:153-167.
70. Greenberg PE, Kessler RC, Birnbaum HG et al. The economic burden of depression in the United States: how did it change between 1990 and 2000? *J Clin Psychiatry.* 2003;64:1465-1475.
71. Ebmeier KP, Donaghey C and Steele JD. Recent developments and current controversies in depression. *Lancet* 2006;367:153-167.
72. Cipriani A, Furukawa TA, Salanti G et al. Comparative efficacy and acceptability of 12 new-generation antidepressants: a multiple-treatments meta-analysis. *Lancet* 2009; e-published January 29, 2009 DOI:10.1016/50140-6736(09)60064-5.
73. Gartlehner G, Gaynes BN, Hansen RA et al. Comparative benefits and harms of second-generation antidepressants: background paper for the American College of Physicians. *Ann Intern Med.* 2008 Nov 18;149(10):734-50.
74. Papakostas GI. Managing partial response or nonresponse: switching, augmentation and combination strategies for major depressive disorder. *J Clin Psychiatry.* 2009;70 Suppl 6:16-25
75. Little A. Treatment-resistant depression. *Am Fam Physician.* 2009 Jul 15;80(2):167-75.
76. Koenig AM, Thase ME. First-line pharmacotherapies for depression-what is the best choice? *Pol Arch Med Wewn.* 2009 Jul-Aug;119(7-8):478-86.
77. Rush A, Roy-Byrne P, Solomon D. Unipolar major depression in adults: Choosing initial treatment. UpToDate. Topic last updated: Nov 28, 2022. Accessed February 23, 2024.

History

Date	Comments
12/13/05	Add to Prescription Drug Section - New Policy—effective January 1, 2006.
08/08/06	Replace Policy - Policy reviewed with literature search by Pharmacy and Therapeutic Committee on July 25, 2006. Policy statement updated with exenatide and thiazolidinediones added as medically necessary; Policy Guidelines and Rationale sections updated; references added.



Date	Comments
05/08/07	Replace Policy - Policy statement for exenatide updated with additional criteria; Policy Guidelines updated to reflect addition to policy statement. Reviewed by P&T on March 27, 2007.
06/12/07	Replace Policy - Policy statement on coverage criteria for exenatide (Byetta), sitagliptin and esomeprazole (Nexium®) expanded; medically necessary indications for 5HT3 antagonists, Actiq and Fentora added to policy statement. Policy Guidelines updated and Rationale updated; references added
12/11/07	Replace Policy - Policy reviewed with literature search by Pharmacy and Therapeutic Committee on May 15, 2007. Policy statement updated to include Pregabalin as either medically necessary or investigational under the criteria. Acyclovir, famciclovir and valacyclovir as medically necessary under criteria. References added.
04/08/08	Replace Policy - Policy updated with literature search by Pharmacy. Policy statement was updated to include fibromyalgia as a medically necessary indication under Pregabalin. References added.
12/16/08	Replace Policy - Policy updated with literature search by Pharmacy. Policy statement updated to include the use of leukotriene modifiers for the treatment of allergic rhinitis refractory to nasal corticosteroids under the medically necessary indication.
02/10/09	New PR Policy PR.5.01.520 - Policy information regarding antidepressants deleted from PR.5.01.605 and addressed in this new policy.
01/12/10	Replace Policy - Policy reviewed with literature search; no change to the policy statement. References added.
05/10/11	Replace Policy - Medically necessary policy statement on branded SSRI, SSNI and second generation antidepressants updated to require trial and failure two generic antidepressants as a condition to be met, where it was previously only one; chronic musculoskeletal pain has been added as an exception to the investigational indications for use of duloxetine (Cymbalta®). Title changed to "Antidepressants: Pharmacy Medical Necessity Criteria for Brands." Reviewed by P&T in March 2011.
09/11/12	Replace policy. Policy updated with literature review. Policy Guidelines section updated to allow for recent availability of generic escitalopram; thus, trial of generic citalopram is no longer a specific requirement for access to Lexapro.
07/08/13	Replace policy. Policy Guidelines updated with Desvenlafaxine, a recently released second-generation SSRI used for treating depression, which may be approved following the failure of two generics, one being venlafaxine. Clarification was added to the policy that it is managed through the member's pharmacy benefit; this is now listed in the header and within the coding section.
12/04/13	Replace policy. Policy section updated with Khedezla™ added to the list of SSRIs which may be approved when criteria are met.



Date	Comments
03/10/14	Replace policy. Cymbalta removed from the scope of policy; prior authorization is no longer required. (NOTE: This is a non-formulary medication; therefore, prior authorization would be required for closed formulary.)
10/13/14	Interim update. Clarification made that policy applies to branded SSRI (selective serotonin reuptake inhibitor) and second generation antidepressants (antidepressants other than tricyclic and MAOI agents) from a single source.
07/14/15	Annual Review. Policy updated with literature review. The following updates were performed: Expanded the indication of Major Depressive Disorder to Depressive Disorders; Expanded the indication of Generalized Anxiety Disorder to Anxiety Disorders; Updated these new expanded indications with separate criteria: For patients with Depressive Disorders, trial and failure of at least two generically available second generation antidepressants, and for patients with Anxiety Disorders, trial and failure of at least two generically available SSRIs, or at least one generically available SSRI and one generically available SNRI.
10/13/15	Interim Update. Requirements of venlafaxine trial for Pristiq, Khedezla or Desvenlafaxine were removed. All brand antidepressants will have same criteria of any 2 generics first. No change to policy statement.
01/01/17	Annual Review, approved December 13, 2016. No published randomized studies were found that would change existing policy statements. Minor correction was made to the 2015 update, i.e., addition of "trial and failure of a generically available SNRI."
12/01/17	Annual Review, approved November 21, 2017. No published randomized studies were found that would change existing policy statements.
11/01/18	Annual Review, approved October 26, 2018. No changes to policy.
01/01/20	Annual Review, approved December 10, 2019. No changes to policy statement.
12/01/20	Annual Review, approved November 3, 2020. No changes to policy statement. Combined depression and anxiety criteria into one table. Added a table documenting that all other uses of brand SSRI, SNRI, and any second- generation antidepressant for this policy are considered not medically necessary. Added a Length of Approval and Documentation Requirements table to policy.
09/01/21	Annual Review, approved August 3, 2021. No changes to policy statement.
11/01/22	Annual Review, approved October 10, 2022. No changes to policy statement. Changed the wording from "patient" to "individual" throughout the policy for standardization.
06/01/23	Annual Review, approved May 22, 2023. No changes to policy statement.
04/01/24	Annual Review, approved March 11, 2024. Added Exxua (gepirone) coverage criteria for the treatment of depression.
01/01/25	Interim Review, approved December 10, 2024. Added Auvelity (dextromethorphan and bupropion) coverage criteria for the treatment of depression. Added to policy the drug



Date	Comments
	names for brand antidepressants. Clarified that the medications listed in this policy are subject to the product's FDA dosage and administration prescribing information.

Disclaimer: This medical policy is a guide in evaluating the medical necessity of a particular service or treatment. The Company adopts policies after careful review of published peer-reviewed scientific literature, national guidelines and local standards of practice. Since medical technology is constantly changing, the Company reserves the right to review and update policies as appropriate. Member contracts differ in their benefits. Always consult the member benefit booklet or contact a member service representative to determine coverage for a specific medical service or supply. CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). ©2025 Premera All Rights Reserved.

Scope: Medical policies are systematically developed guidelines that serve as a resource for Company staff when determining coverage for specific medical procedures, drugs or devices. Coverage for medical services is subject to the limits and conditions of the member benefit plan. Members and their providers should consult the member benefit booklet or contact a customer service representative to determine whether there are any benefit limitations applicable to this service or supply. This medical policy does not apply to Medicare Advantage.

